

D-branes and Deep Learning

Theoretical and Computational Aspects in String Theory

Riccardo Finotello

Scuola di Dottorato in Fisica e Astrofisica

Università degli Studi di Torino
and
I.N.F.N. – sezione di Torino

18th December 2020



Contents

Conformal Symmetry and Geometry of the Worldsheet

Preliminary Concepts and Tools

D-branes Intersecting at Angles

Fermions and Point-like Defect CFT

Cosmological Backgrounds and Divergences

Orbifolds and Cosmological Toy Models

Null Boost Orbifold

Deep Learning the Geometry of String Theory

Machine Learning and Deep Learning

Machine Learning for String Theory

AI Implementations for Geometry and Strings

Contents

Conformal Symmetry and Geometry of the Worldsheet

Preliminary Concepts and Tools

D-branes Intersecting at Angles

Fermions and Point-like Defect CFT

Cosmological Backgrounds and Divergences

Orbifolds and Cosmological Toy Models

Null Boost Orbifold

Deep Learning the Geometry of String Theory

Machine Learning and Deep Learning

Machine Learning for String Theory

AI Implementations for Geometry and Strings

Action Principle and Conformal Symmetry

Polyakov's Action

$$S_P[\gamma, X, \psi] = -\frac{1}{4\pi} \int_{-\infty}^{+\infty} d\tau \int_0^\ell d\sigma \sqrt{-\det \gamma} \gamma^{\alpha\beta} \left(\frac{2}{\alpha'} \partial_\alpha X^\mu \partial_\beta X^\nu + \psi^\mu \rho_\alpha \partial_\beta \psi^\nu \right) \eta_{\mu\nu}$$

Action Principle and Conformal Symmetry

Polyakov's Action

$$S_P[\gamma, X, \psi] = -\frac{1}{4\pi} \int_{-\infty}^{+\infty} d\tau \int_0^\ell d\sigma \sqrt{-\det \gamma} \gamma^{\alpha\beta} \left(\frac{2}{\alpha'} \partial_\alpha X^\mu \partial_\beta X^\nu + \psi^\mu \rho_\alpha \partial_\beta \psi^\nu \right) \eta_{\mu\nu}$$

Symmetries:

Poincaré transf.: $X'^\mu = \Lambda^\mu_\nu X^\nu + c^\mu$
2D diff.: $\gamma'_{\alpha\beta} = (J^{-1})_{\alpha\beta}^{\lambda\rho} \gamma_{\lambda\rho}$
Weyl transf.: $\gamma'_{\alpha\beta} = e^{2\omega} \gamma_{\alpha\beta}$

Conformal symmetry:

vanishing stress-energy tensor: $\mathcal{T}_{\alpha\beta} = 0$
traceless stress-energy tensor: $\text{tr } \mathcal{T} = 0$
conformal gauge: $\gamma_{\alpha\beta} = e^\Phi \eta_{\alpha\beta}$

Action Principle and Conformal Symmetry

Superstrings in D dimensions $\longrightarrow$ *Virasoro algebra* (central extension of de Witt's algebra):

$$\mathcal{T}(z) = -\frac{1}{\alpha'} \partial X(z) \cdot \partial X(z) - \frac{1}{2} \psi(z) \cdot \partial \psi(z) \quad \Rightarrow \quad c = \frac{3}{2} D$$

Action Principle and Conformal Symmetry

Superstrings in D dimensions $\longrightarrow$ *Virasoro algebra* (central extension of de Witt's algebra):

$$\mathcal{T}(z) = -\frac{1}{\alpha'} \partial X(z) \cdot \partial X(z) - \frac{1}{2} \psi(z) \cdot \partial \psi(z) \quad \Rightarrow \quad c = \frac{3}{2} D$$

$(\lambda, 0) / (1 - \lambda, 0)$ Ghost System

Introduce anti-commuting (b, c) and commuting (β, γ) conformal fields:

$$S_{\text{ghost}}[b, c, \beta, \gamma] = \frac{1}{2\pi} \iint dz d\bar{z} (b(z) \bar{\partial} c(z) + \beta(z) \bar{\partial} \gamma(z))$$

where $\lambda_b = 2$ and $\lambda_c = -1$, and $\lambda_\beta = \frac{3}{2}$ and $\lambda_\gamma = -\frac{1}{2}$.

[Friedan, Martinec, Shenker (1986)]

Action Principle and Conformal Symmetry

Superstrings in D dimensions $\longrightarrow$ *Virasoro algebra* (central extension of de Witt's algebra):

$$\mathcal{T}(z) = -\frac{1}{\alpha'} \partial X(z) \cdot \partial X(z) - \frac{1}{2} \psi(z) \cdot \partial \psi(z) \quad \Rightarrow \quad c = \frac{3}{2} D$$

$(\lambda, 0) / (1 - \lambda, 0)$ Ghost System

Introduce anti-commuting (b, c) and commuting (β, γ) conformal fields:

$$S_{\text{ghost}}[b, c, \beta, \gamma] = \frac{1}{2\pi} \iint dz d\bar{z} (b(z) \bar{\partial} c(z) + \beta(z) \bar{\partial} \gamma(z))$$

where $\lambda_b = 2$ and $\lambda_c = -1$, and $\lambda_\beta = \frac{3}{2}$ and $\lambda_\gamma = -\frac{1}{2}$.

[Friedan, Martinec, Shenker (1986)]

Consequence:


$$c_{\text{full}} = c + c_{\text{ghost}} = 0 \quad \Leftrightarrow \quad D = 10.$$

Extra Dimensions and Compactification

Compactification

$$\mathcal{M}^{1,9} = \mathcal{M}^{1,3} \otimes \mathcal{X}_6$$

- $\mathcal{X}_6$ is a **compact** manifold
- $N = 1$ **supersymmetry** preserved in 4D
- contains algebra of $SU(3) \otimes SU(2) \otimes U(1)$




[code in Hanson (1994)]

Extra Dimensions and Compactification

Compactification

$$\mathcal{M}^{1,9} = \mathcal{M}^{1,3} \otimes \mathcal{X}_6$$

- $\mathcal{X}_6$ is a **compact** manifold
- $N = 1$ **supersymmetry** preserved in 4D
- contains algebra of $SU(3) \otimes SU(2) \otimes U(1)$



[code in Hanson (1994)]

Calabi–Yau manifolds (M, g) such that:

- $\dim_{\mathbb{C}} M = m$
- $\text{Hol}(g) \subseteq SU(m)$
- $\text{Ric}(g) \equiv 0$ (equiv. $c_1(M) \equiv 0$)


[Calabi (1957), Yau (1977), Candelas et al. (1985)]

Extra Dimensions and Compactification

Compactification

$$\mathcal{M}^{1,9} = \mathcal{M}^{1,3} \otimes \mathcal{X}_6$$

- $\mathcal{X}_6$ is a **compact** manifold
- $N = 1$ **supersymmetry** preserved in 4D
- contains algebra of $SU(3) \otimes SU(2) \otimes U(1)$



[code in Hanson (1994)]

Calabi–Yau manifolds (M, g) such that:

- $\dim_{\mathbb{C}} M = m$
- $\text{Hol}(g) \subseteq SU(m)$
- $\text{Ric}(g) \equiv 0$ (equiv. $c_1(M) \equiv 0$)

[Calabi (1957), Yau (1977), Candelas et al. (1985)]

Characterised by **Hodge numbers**

$$h^{r,s} = \dim_{\mathbb{C}} H_{\bar{\partial}}^{r,s}(M, \mathbb{C})$$

(no. of harmonic (r, s) -forms).

D-branes and Open Strings

Polyakov's action naturally introduces **Neumann b.c.**:

$$\partial_\sigma X(\tau, \sigma) \Big|_{\sigma=0}^{\sigma=\ell} = 0$$

satisfied by **open and closed strings** in D dim. s.t. $\square X = 0 \Rightarrow X(z, \bar{z}) = X(z) + \overline{X}(\bar{z})$.

D-branes and Open Strings

Polyakov's action naturally introduces **Neumann b.c.**:

$$\partial_\sigma X(\tau, \sigma) \Big|_{\sigma=0}^{\sigma=\ell} = 0$$

satisfied by **open and closed strings** in D dim. s.t. $\square X = 0 \Rightarrow X(z, \bar{z}) = X(z) + \bar{X}(\bar{z})$.

T-duality

Consider **closed strings** on $\mathcal{M}^{1,D-1} = \mathcal{M}^{1,D-2} \otimes S^1(R)$:

$$\begin{aligned} \begin{cases} \alpha_0^{D-1} &= \frac{1}{\sqrt{2\alpha'}} \left(n \frac{\alpha'}{R} + mR \right) \\ \tilde{\alpha}_0^{D-1} &= \frac{1}{\sqrt{2\alpha'}} \left(n \frac{\alpha'}{R} - mR \right) \end{cases} \Rightarrow M^2 = -p^\mu p_\mu = \frac{2}{\alpha'} (\alpha_0^{D-1})^2 + \frac{4}{\alpha'} (N + a) \\ &= \frac{2}{\alpha'} (\tilde{\alpha}_0^{D-1})^2 + \frac{4}{\alpha'} (\tilde{N} + a) \end{aligned}$$

D-branes and Open Strings

Polyakov's action naturally introduces **Neumann b.c.**:

$$\partial_\sigma X(\tau, \sigma) \Big|_{\sigma=0}^{\sigma=\ell} = 0$$

satisfied by **open and closed strings** in D dim. s.t. $\square X = 0 \Rightarrow X(z, \bar{z}) = X(z) + \bar{X}(\bar{z})$.

T-duality

Dirichlet b.c. consequence of **T-duality** on p directions:

$$\bar{X}(z) \mapsto -\bar{X}(z) \quad \Rightarrow \quad \partial_\sigma X^i(\tau, \sigma) \Big|_{\sigma=0}^{\sigma=\ell} = 0 \quad \xrightarrow{T\text{-duality}} \quad \partial_\tau \tilde{X}^i(\tau, \sigma) \Big|_{\sigma=0}^{\sigma=\ell} = 0$$

thus **open strings** can be **constrained** to $D(D - p - 1)$ -branes.

[Polchinski (1995, 1996)]

D-branes and Open Strings


Introducing Dp -branes breaks $\text{ISO}(1, D-1) \rightarrow \text{ISO}(1, p) \otimes \text{SO}(D-1-p)$:

$$\mathcal{A}^\mu \rightarrow (\mathcal{A}^A, \mathcal{A}^a) \quad \Rightarrow \quad \text{U}(1) \text{ theory in } p+1 \text{ dimensions (and scalars)}$$

D-branes and Open Strings

Introducing Dp -branes breaks $\text{ISO}(1, D-1) \rightarrow \text{ISO}(1, p) \otimes \text{SO}(D-1-p)$:

$$\mathcal{A}^\mu \rightarrow (\mathcal{A}^A, \mathcal{A}^a) \Rightarrow \text{U}(1) \text{ theory in } p+1 \text{ dimensions (and scalars)}$$




[Chan, Paton (1969)]

$$|n; r\rangle = \sum_{i,j=1}^N |n; i, j\rangle \lambda^r_{ij} \Rightarrow \text{U}(N)$$

D-branes and Open Strings


Introducing Dp -branes breaks $\text{ISO}(1, D-1) \rightarrow \text{ISO}(1, p) \otimes \text{SO}(D-1-p)$:

$$\mathcal{A}^\mu \rightarrow (\mathcal{A}^A, \mathcal{A}^a) \Rightarrow \text{U}(1) \text{ theory in } p+1 \text{ dimensions (and scalars)}$$



[Chan, Paton (1969)]

$$|n; r\rangle = \sum_{i,j=1}^N |n; i, j\rangle \lambda^r_{ij} \Rightarrow \text{U}(N)$$



Intersecting D-branes

Consider 3 intersecting *D6*-branes filling $\mathcal{M}^{1,3}$ and embedded in $\mathbb{R}^6$

Twist Fields Correlators

$$\left\langle \prod_{t=1}^{N_B} \sigma_{M_{(t)}}(x_{(t)}) \right\rangle = \mathcal{N} \left(\{x_{(t)}, M_{(t)}\}_{1 \leq t \leq N_B} \right) e^{-S_{E(\text{cl})} \left(\{x_{(t)}, M_{(t)}\}_{1 \leq t \leq N_B} \right)}$$

Intersecting D-branes

Consider 3 intersecting *D6*-branes filling $\mathcal{M}^{1,3}$ and embedded in $\mathbb{R}^6$

Twist Fields Correlators


$$\left\langle \prod_{t=1}^{N_B} \sigma_{M_{(t)}}(x_{(t)}) \right\rangle = \mathcal{N} \left(\{x_{(t)}, M_{(t)}\}_{1 \leq t \leq N_B} \right) e^{-S_E(\text{cl}) \left(\{x_{(t)}, M_{(t)}\}_{1 \leq t \leq N_B} \right)}$$

Intersecting D-branes

Consider 3 intersecting $D6$ -branes filling $\mathcal{M}^{1,3}$ and embedded in $\mathbb{R}^6$

Twist Fields Correlators

$$\left\langle \prod_{t=1}^{N_B} \sigma_{M(t)}(x(t)) \right\rangle = \mathcal{N} \left(\{x(t), M(t)\}_{1 \leq t \leq N_B} \right) e^{-S_E(\text{cl}) \left(\{x(t), M(t)\}_{1 \leq t \leq N_B} \right)}$$




D-branes in **factorised** internal space:

- **embedded as lines** in $\mathbb{R}^2 \times \mathbb{R}^2 \times \mathbb{R}^2$
- **relative rotations** are $SO(2) \simeq U(1)$ elements
- $S_E^{(\text{cl})} \left(\{x(t), M(t)\}_{1 \leq t \leq N_B} \right) \sim \text{Area} \left(\{f(t), R(t)\}_{1 \leq t \leq N_B} \right)$

[Cremades, Ibanez, Marchesano (2003); Pesando (2012)]

SO(4) Rotations

Consider $\mathbb{R}^4 \times \mathbb{R}^2$ (focus on $\mathbb{R}^4$):



$$(X_{(t)})^I = (R_{(t)})^I_J X^J - g_{(t)}^I \in \mathbb{R}^4$$

where


$$R_{(t)} \in \frac{\text{SO}(4)}{\text{S}(\text{O}(2) \times \text{O}(2))}$$

that is

$$[R_{(t)}] = \{R_{(t)} \sim \mathcal{O}_{(t)} R_{(t)}\}$$


Boundary Conditions and Open Strings

- $u = x + iy = e^{\tau_e + i\sigma}$ and $\bar{u} = u^*$
- $x_{(t)} < x_{(t-1)}$ **worldsheet intersection points**



Boundary Conditions and Open Strings

- $u = x + iy = e^{\tau_e + i\sigma}$ and $\bar{u} = u^*$
- $x_{(t)} < x_{(t-1)}$ **worldsheet intersection points**



Branch Cuts and Discontinuities for $x \in D_{(t)}$

$$\begin{cases} \partial_u X(x + i0^+) = U_{(t)} \cdot \partial_{\bar{u}} \bar{X}(x - i0^+) = \left[R_{(t)}^{-1} \cdot (\sigma_3 \otimes \mathbb{1}_2) \cdot R_{(t)} \right] \cdot \partial_{\bar{u}} \bar{X}(x - i0^+) \\ X(x_{(t)}, x_{(t)}) = f_{(t)} \end{cases}$$

Doubling Trick and Spinor Representation

Doubling Trick

$$\partial_z \mathcal{X}(z) = \begin{cases} \partial_u X(u) & \text{if } z \in \mathcal{H}_{>}^{(\bar{t})} \\ U_{(\bar{t})} \partial_{\bar{u}} \bar{X}(\bar{u}) & \text{if } z \in \mathcal{H}_{<}^{(\bar{t})} \end{cases} \Rightarrow \begin{aligned} \partial_z \mathcal{X}(x_{(t)} + e^{2\pi i} \delta_+) &= \mathcal{U}_{(t, t+1)} \partial_z \mathcal{X}(x_{(t)} + \delta_+), \\ \partial_z \mathcal{X}(x_{(t)} + e^{2\pi i} \delta_-) &= \tilde{\mathcal{U}}_{(t, t+1)} \partial_z \mathcal{X}(x_{(t)} + \delta_-), \end{aligned}$$

where $\mathcal{H}_{\gtrless}^{(t)} = \{z \in \mathbb{C} \mid \text{Im } z \gtrless 0 \text{ or } z \in D_{(t)}\}$ and $\delta_{\pm} = \eta \pm i0^+$.

Doubling Trick and Spinor Representation

Doubling Trick

$$\partial_z \mathcal{X}(z) = \begin{cases} \partial_u X(u) & \text{if } z \in \mathcal{H}_{>}^{(\bar{t})} \\ U_{(\bar{t})} \partial_{\bar{u}} \bar{X}(\bar{u}) & \text{if } z \in \mathcal{H}_{<}^{(\bar{t})} \end{cases} \Rightarrow \begin{aligned} \partial_z \mathcal{X}(x_{(t)} + e^{2\pi i} \delta_+) &= \mathcal{U}_{(t, t+1)} \partial_z \mathcal{X}(x_{(t)} + \delta_+), \\ \partial_z \mathcal{X}(x_{(t)} + e^{2\pi i} \delta_-) &= \tilde{\mathcal{U}}_{(t, t+1)} \partial_z \mathcal{X}(x_{(t)} + \delta_-), \end{aligned}$$

where $\mathcal{H}_{\gtrless}^{(t)} = \{z \in \mathbb{C} \mid \text{Im } z \gtrless 0 \text{ or } z \in D_{(t)}\}$ and $\delta_{\pm} = \eta \pm i0^+$.

Doubling Trick and Spinor Representation

Doubling Trick

$$\partial_z \mathcal{X}(z) = \begin{cases} \partial_u X(u) & \text{if } z \in \mathcal{H}_{>}^{(\bar{t})} \\ U_{(\bar{t})} \partial_{\bar{u}} \bar{X}(\bar{u}) & \text{if } z \in \mathcal{H}_{<}^{(\bar{t})} \end{cases} \Rightarrow \begin{aligned} \partial_z \mathcal{X}(x_{(t)} + e^{2\pi i} \delta_+) &= \mathcal{U}_{(t, t+1)} \partial_z \mathcal{X}(x_{(t)} + \delta_+), \\ \partial_z \mathcal{X}(x_{(t)} + e^{2\pi i} \delta_-) &= \tilde{\mathcal{U}}_{(t, t+1)} \partial_z \mathcal{X}(x_{(t)} + \delta_-), \end{aligned}$$

where $\mathcal{H}_{\gtrless}^{(t)} = \{z \in \mathbb{C} \mid \text{Im } z \gtrless 0 \text{ or } z \in D_{(t)}\}$ and $\delta_{\pm} = \eta \pm i0^+$.


Use Pauli matrices $\tau = (i \mathbb{1}_2, \vec{\sigma})$:

$$\partial_z \mathcal{X}_{(s)}(z) = \partial_z \mathcal{X}'(z) \tau_I \Rightarrow \partial_z \mathcal{X}(x_{(t)} + e^{2\pi i} \delta_{\pm}) = \overset{(\sim)}{\mathcal{L}}_{(t, t+1)} \partial_z \mathcal{X}(x_{(t)} + \delta_{\pm}) \overset{(\sim)}{\mathcal{R}}_{(t, t+1)}$$

where

$$\overset{(\sim)}{\mathcal{L}}_{(t, t+1)} \in \text{SU}(2)_L \quad \text{and} \quad \overset{(\sim)}{\mathcal{R}}_{(t, t+1)} \in \text{SU}(2)_R$$


Hypergeometric Basis



Sum over all contributions:

$$\partial_z \mathcal{X}(z) = \frac{\partial \omega_z}{\partial z} \sum_{l, r=-\infty}^{+\infty} c_{lr} (-\omega_z)^{A_{lr}} (1 - \omega_z)^{B_{lr}} \times B_{0,l}^{(L)}(\omega_z) \left(B_{0,r}^{(R)}(\omega_z) \right)^T$$

Hypergeometric Basis



Sum over all contributions:

$$\partial_z \mathcal{X}(z) = \frac{\partial \omega_z}{\partial z} \sum_{l, r=-\infty}^{+\infty} c_{lr} (-\omega_z)^{A_{lr}} (1 - \omega_z)^{B_{lr}} \times B_{0,l}^{(L)}(\omega_z) \left(B_{0,r}^{(R)}(\omega_z) \right)^T$$

Basis of Solutions

$$B_{0,n}(\omega_z) = \begin{pmatrix} 1 & 0 \\ 0 & K_n \end{pmatrix} \begin{pmatrix} \frac{1}{\Gamma(c_n)} {}_2F_1(a_n, b_n; c_n; \omega_z) \\ \frac{(-\omega_z)^{1-c_n}}{\Gamma(2-c_n)} {}_2F_1(a_n + 1 - c_n, b_n + 1 - c_n; 2 - c_n; \omega_z) \end{pmatrix}$$

The Solution

Sequence of the operations:


1. rotation matrix = monodromy matrix
2. contiguity relations $\Rightarrow$ independent hypergeometrics
3. finite action $\Rightarrow$ 2 solutions (no. of d.o.f. is correctly saturated)
4. boundary conditions $\Rightarrow$ fix free constants c_{lr}

The Solution

Sequence of the operations:

1. rotation matrix = monodromy matrix
2. contiguity relations $\Rightarrow$ independent hypergeometrics
3. finite action $\Rightarrow$ 2 solutions (no. of d.o.f. is correctly saturated)
4. boundary conditions $\Rightarrow$ fix free constants c_{lr}

Physical Interpretation




$$\begin{aligned}
 2\pi\alpha' S_{\mathbb{R}^4} \Big|_{\text{on-shell}} &= \sum_{t=1}^3 \left(\frac{1}{2} \left| g_{(t)}^\perp \right| \left| f_{(t-1)} - f_{(t)} \right| \right) \\
 &= \text{Area} \left(\{ f_{(t)} \}_{1 \leq t \leq N_B} \right)
 \end{aligned}$$

The Solution

Sequence of the operations:

1. rotation matrix = monodromy matrix
2. contiguity relations $\Rightarrow$ independent hypergeometrics
3. finite action $\Rightarrow$ 2 solutions (no. of d.o.f. is correctly saturated)
4. boundary conditions $\Rightarrow$ fix free constants c_{lr}


Physical Interpretation



- strings no longer confined to plane
- strings form a *small bump* from the D-brane
- classical action **larger** than factorised case

[RF, Pesando (2019)]


Fermions on the Strip



Action of Boundary Changing Operators

$$\begin{cases} \psi_-^i(\tau, 0) &= (R_{(t)})^I{}_J \psi_+^J(\tau, 0) \quad \text{for } \tau \in (\hat{\tau}_{(t)}, \hat{\tau}_{(t-1)}) \\ \psi_-^I(\tau, \pi) &= -\psi_+^I(\tau, \pi) \quad \text{for } \tau \in \mathbb{R} \end{cases}$$

Fermions on the Strip



Action of Boundary Changing Operators

$$\begin{cases} \psi_-^i(\tau, 0) &= (R_{(t)})^I{}_J \psi_+^J(\tau, 0) \quad \text{for } \tau \in (\hat{\tau}_{(t)}, \hat{\tau}_{(t-1)}) \\ \psi_-^I(\tau, \pi) &= -\psi_+^I(\tau, \pi) \quad \text{for } \tau \in \mathbb{R} \end{cases}$$

$$\mathcal{T}_{\pm\pm}(\xi_{\pm}) = -i \frac{T}{4} \psi_{\pm, I}^*(\xi_{\pm}) \overset{\leftrightarrow}{\partial} \psi_{\pm}^I(\xi_{\pm}) \Rightarrow \begin{cases} \dot{H}(\tau) &= 0 \\ \dot{P}(\tau) &\neq 0 \end{cases} \Leftrightarrow \tau \in (\tau_{(t)}, \tau_{(t-1)})$$

Conserved Product and Operators

Expand on a **basis of solutions**

$$\psi_{\pm}(\xi_{\pm}) = \sum_{n=-\infty}^{+\infty} b_n \psi_n(\xi_{\pm}) \quad \Rightarrow \quad \Psi(z) = \begin{cases} \psi_{E,+}(u) & \text{if } z \in \mathcal{H}_{>}^{(\bar{t})} \\ \psi_{E,-}(u) & \text{if } z \in \mathcal{H}_{<}^{(\bar{t})} \end{cases}$$

Conserved Product and Operators

Expand on a **basis of solutions**

$$\psi_{\pm}(\xi_{\pm}) = \sum_{n=-\infty}^{+\infty} b_n \psi_n(\xi_{\pm}) \quad \Rightarrow \quad \Psi(z) = \begin{cases} \psi_{E,+}(u) & \text{if } z \in \mathcal{H}_{>}^{(\bar{t})} \\ \psi_{E,-}(u) & \text{if } z \in \mathcal{H}_{<}^{(\bar{t})} \end{cases}$$

Conserved Product and Dual Basis

$$\langle\langle {}^*\psi_n, \psi_m \rangle\rangle = 2\pi\mathcal{N} \oint \frac{dz}{2\pi i} {}^*\Psi_n {}^*\Psi_m = \delta_{n,m} \quad \Rightarrow \quad \langle\langle {}^*\Psi_n^{(*)}, \Psi^{(*)} \rangle\rangle = b_n^{(\dagger)}$$

Conserved Product and Operators

Expand on a **basis of solutions**

$$\psi_{\pm}(\xi_{\pm}) = \sum_{n=-\infty}^{+\infty} b_n \psi_n(\xi_{\pm}) \quad \Rightarrow \quad \Psi(z) = \begin{cases} \psi_{E,+}(u) & \text{if } z \in \mathcal{H}_{>}^{(\bar{t})} \\ \psi_{E,-}(u) & \text{if } z \in \mathcal{H}_{<}^{(\bar{t})} \end{cases}$$

Conserved Product and Dual Basis

$$\langle\langle {}^*\psi_n, \psi_m \rangle\rangle = 2\pi\mathcal{N} \oint \frac{dz}{2\pi i} {}^*\Psi_n {}^*\Psi_m = \delta_{n,m} \quad \Rightarrow \quad \langle\langle {}^*\Psi_n^{(*)}, \Psi^{(*)} \rangle\rangle = b_n^{(\dagger)}$$

Derive the **algebra of operators**:

$$[b_n, b_m^{\dagger}]_+ = \frac{2\mathcal{N}}{T} \langle\langle {}^*\Psi_n^*, \Psi_m^* \rangle\rangle$$

Twisted Complex Fermions

Consider $R_{(t)} = e^{i\pi\alpha_{(t)}} \in U(1)$:

$$\Psi(x_{(t)} + e^{2\pi i} \delta) = e^{i\pi\epsilon_{(t)}} \Psi(x_{(t)} + \delta)$$

where

$$\epsilon_{(t)} = \alpha_{(t+1)} - \alpha_{(t)} + \theta(\alpha_{(t)} - \alpha_{(t+1)} - 1) - \theta(\alpha_{(t+1)} - \alpha_{(t)} - 1)$$

Twisted Complex Fermions

Consider $R_{(t)} = e^{i\pi\alpha_{(t)}} \in U(1)$:

$$\Psi(x_{(t)} + e^{2\pi i}\delta) = e^{i\pi\epsilon_{(t)}} \Psi(x_{(t)} + \delta)$$

where

$$\epsilon_{(t)} = \alpha_{(t+1)} - \alpha_{(t)} + \theta(\alpha_{(t)} - \alpha_{(t+1)} - 1) - \theta(\alpha_{(t+1)} - \alpha_{(t)} - 1)$$

Basis of Solutions

$$\Psi_n(z; \{x_{(t)}\}) = \mathcal{N}_\Psi z^{-n} \prod_{t=1}^N \left(1 - \frac{z}{x_{(t)}}\right)^{n_{(t)} + \frac{\epsilon_{(t)}}{2}}$$

$$^*\Psi_n(z; \{x_{(t)}\}) = \frac{1}{2\pi\mathcal{N}\mathcal{N}_\Psi} z^{n-1} \prod_{t=1}^N \left(1 - \frac{z}{x_{(t)}}\right)^{-\tilde{n}_{(t)} + \frac{\epsilon_{(t)}}{2}}$$

Vacua

Define the **vacuum** with respect to b_n :

$$b_n \left| \{x_{(t)}\} \right\rangle = 0 \quad \text{for } n \geq 1$$

$$b_n \left| \tilde{0} \right\rangle = 0 \quad \text{for } n \geq n_{(t)} + \frac{\epsilon_{(t)}}{2} + \frac{1}{2}$$

Vacua


Define the **vacuum** with respect to b_n :

$$b_n \left| \{x_{(t)}\} \right\rangle = 0 \quad \text{for } n \geq 1$$

$$b_n \left| \tilde{0} \right\rangle = 0 \quad \text{for } n \geq n_{(t)} + \frac{\epsilon_{(t)}}{2} + \frac{1}{2}$$

Theories are subject to **consistency conditions**:

$$\langle \{x_{(t)}\} | \{x_{(t)}\} \rangle = 1 \quad \Rightarrow \quad L = n_{(t)} + \tilde{n}_{(t)}$$



Vacua


Define the **vacuum** with respect to b_n :

$$b_n \left| \{x_{(t)}\} \right\rangle = 0 \quad \text{for } n \geq 1$$

$$b_n \left| \tilde{0} \right\rangle = 0 \quad \text{for } n \geq n_{(t)} + \frac{\epsilon_{(t)}}{2} + \frac{1}{2}$$

Theories are subject to **consistency conditions**:

$$\langle \{x_{(t)}\} | \{x_{(t)}\} \rangle = 1 \quad \Rightarrow \quad L = n_{(t)} + \tilde{n}_{(t)} = 0$$



Stress-energy Tensor and CFT Approach

Compute the OPEs leading to the **time dependent stress-energy tensor**:

$$\mathcal{T}(z) = \frac{\pi T}{2} \mathcal{N}_{\Psi}^2 \sum_{n, m=-\infty}^{+\infty} : b_n b_m^* : z^{-n-m} \left[\frac{m-n}{2} + 2 \sum_{t=1}^N \frac{n_{(t)} + \frac{\epsilon_{(t)}}{2}}{z - x_{(t)}} \right] + \frac{1}{2} \left(\sum_{t=1}^N \frac{n_{(t)} + \frac{\epsilon_{(t)}}{2}}{z - x_{(t)}} \right)^2$$

[RF, Pesando (2019)]

Stress-energy Tensor and CFT Approach

Compute the OPEs leading to the **time dependent stress-energy tensor**:

$$\mathcal{T}(z) = \frac{\pi T}{2} \mathcal{N}_{\Psi}^2 \sum_{n, m=-\infty}^{+\infty} : b_n b_m^* : z^{-n-m} \left[\frac{m-n}{2} + 2 \sum_{t=1}^N \frac{n(t) + \frac{\epsilon(t)}{2}}{z - x(t)} \right] + \frac{1}{2} \left(\sum_{t=1}^N \frac{n(t) + \frac{\epsilon(t)}{2}}{z - x(t)} \right)^2$$

[RF, Pesando (2019)]

Invariant Vacuum and Spin Fields

$$|\{x(t)\}\rangle = \mathcal{N}(\{x(t)\}) \mathcal{R} \left[\prod_{t=1}^M S_{(t)}(x(t)) \right] |0\rangle_{\text{SL}_2(\mathbb{R})}$$

Spin Fields Amplitudes

Equivalence with Bosonization

$$\partial_{x_{(t)}} \ln \langle \{x_{(t)}\} | \{x_{(t)}\} \rangle = \oint_{x_{(t)}} \frac{dz}{2\pi i} \frac{\langle \{x_{(t)}\} | \mathcal{T}(z) | \{x_{(t)}\} \rangle}{\langle \{x_{(t)}\} | \{x_{(t)}\} \rangle}$$

$$\Rightarrow \langle \{x_{(t)}\} | \{x_{(t)}\} \rangle = \mathcal{N}(\{\epsilon_{(t)}\}) \prod_{\substack{t=1 \\ t>u}}^N (x_{(u)} - x_{(t)})^{\left(n_{(u)} + \frac{\epsilon_{(u)}}{2}\right) \left(n_{(t)} + \frac{\epsilon_{(t)}}{2}\right)}$$

Spin Fields Amplitudes

Equivalence with Bosonization

$$\partial_{x_{(t)}} \ln \langle \{x_{(t)}\} | \{x_{(t)}\} \rangle = \oint_{x_{(t)}} \frac{dz}{2\pi i} \frac{\langle \{x_{(t)}\} | \mathcal{T}(z) | \{x_{(t)}\} \rangle}{\langle \{x_{(t)}\} | \{x_{(t)}\} \rangle}$$

$$\Rightarrow \langle \{x_{(t)}\} | \{x_{(t)}\} \rangle = \mathcal{N}(\{\epsilon_{(t)}\}) \prod_{\substack{t=1 \\ t > u}}^N (x_{(u)} - x_{(t)})^{\left(n_{(u)} + \frac{\epsilon_{(u)}}{2}\right) \left(n_{(t)} + \frac{\epsilon_{(t)}}{2}\right)}$$

- (semi-)phenomenological models involve **twist and spin** fields and **open strings**
- framework for **bosonic** open strings with **intersecting D-branes**
- **spin fields** as **boundary changing operators** (hidden in **defects**)
- framework for amplitudes (extension to (non) Abelian twist/spin fields?)

Contents

Conformal Symmetry and Geometry of the Worldsheet

Preliminary Concepts and Tools

D-branes Intersecting at Angles

Fermions and Point-like Defect CFT

Cosmological Backgrounds and Divergences

Orbifolds and Cosmological Toy Models

Null Boost Orbifold

Deep Learning the Geometry of String Theory

Machine Learning and Deep Learning

Machine Learning for String Theory


AI Implementations for Geometry and Strings

A Few Words on a Theory of Everything

string theory = theory of everything = **nuclear forces** + **gravity**

A Few Words on a Theory of Everything

string theory = theory of everything = **nuclear forces** + **gravity**




From the phenomenological point of view:

- cosmological implications

A Few Words on a Theory of Everything

string theory = theory of everything = **nuclear forces** + **gravity**




From the phenomenological point of view:

- cosmological implications
- Big Bang(-like) singularities

A Few Words on a Theory of Everything

string theory = theory of everything = **nuclear forces** + **gravity**




From the phenomenological point of view:

- cosmological implications
- Big Bang(-like) singularities
- toy models of **space-like singularities**

A Few Words on a Theory of Everything

string theory = theory of everything = **nuclear forces** + **gravity**



From the phenomenological point of view:

- cosmological implications
- Big Bang(-like) singularities
- toy models of **space-like singularities**



time-dependent orbifold models

[Craps, Kutasov, Rajesh (2002); Liu, Moore, Seiberg (2002)]

Cosmological Singularities

Use **time-dependent orbifolds** to model **space-like singularities**:

divergent closed string amplitudes $\Rightarrow$ gravitational backreaction?

Cosmological Singularities

Use **time-dependent orbifolds** to model **space-like singularities**:

divergent closed string amplitudes $\Rightarrow$ gravitational backreaction?

Divergences

Even in simple models (e.g. NBO, more on this later) the 4 tachyons amplitude is divergent **in the open sector at tree level**:

$$A_4 \sim \int_{q \sim \infty} \frac{dq}{|q|} \mathcal{A}(q)$$

where

$$\mathcal{A}_{\text{closed}}(q) \sim q^{4-\alpha' \|\vec{p}_\perp\|^2} \quad \text{and} \quad \mathcal{A}_{\text{open}}(q) \sim q^{1-\alpha' \|\vec{p}_\perp\|^2} \text{tr}([T_1, T_2]_+ [T_3, T_4]_+)$$

Null Boost Orbifold

Start from $(x^+, x^-, x^2, \vec{x}) \in \mathcal{M}^{1,D-1}$:

$$\begin{cases} u &= x^- \\ z &= \frac{x^2}{\Delta x^-} \\ v &= x^+ - \frac{1}{2} \frac{(x^2)^2}{x^-} \end{cases} \quad \Rightarrow \quad ds^2 = -2 du dv + (\Delta u)^2 dz^2 + \delta_{ij} dx^i dx^j$$

Null Boost Orbifold

Start from $(x^+, x^-, x^2, \vec{x}) \in \mathcal{M}^{1,D-1}$:

$$\begin{cases} u &= x^- \\ z &= \frac{x^2}{\Delta x^-} \\ v &= x^+ - \frac{1}{2} \frac{(x^2)^2}{x^-} \end{cases} \quad \Rightarrow \quad ds^2 = -2 du dv + (\Delta u)^2 dz^2 + \delta_{ij} dx^i dx^j$$

Killing Vector and Null Boost Orbifold

$$\kappa = -i(2\pi\Delta)J_{+2} = 2\pi\partial_z \quad \Rightarrow \quad z \sim z + 2\pi n$$

Null Boost Orbifold

Start from $(x^+, x^-, x^2, \vec{x}) \in \mathcal{M}^{1,D-1}$:

$$\begin{cases} u &= x^- \\ z &= \frac{x^2}{\Delta x^-} \\ v &= x^+ - \frac{1}{2} \frac{(x^2)^2}{x^-} \end{cases} \Rightarrow ds^2 = -2 du dv + (\Delta u)^2 dz^2 + \delta_{ij} dx^i dx^j$$

Killing Vector and Null Boost Orbifold

$$\kappa = -i(2\pi\Delta)J_{+2} = 2\pi\partial_z \Rightarrow z \sim z + 2\pi n$$

Scalars on NBO:

$$\Phi_{\{k_+, l, \vec{k}, r\}}(u, v, z, \vec{x}) = e^{i(k_+ v + l z + \vec{k} \cdot \vec{x})} \tilde{\Phi}_{\{k_+, l, \vec{k}, r\}}(u) = \frac{e^{i(k_+ v + l z + \vec{k} \cdot \vec{x})}}{\sqrt{(2\pi)^D |2\Delta k_+ u|}} e^{-i \frac{l^2}{2\Delta^2 k_+} \frac{1}{u} + i \frac{\|\vec{k}\|^2 + r}{2k_+} u}$$

Scalar QED Interactions

Scalar-photon interactions:

$$S_{\text{sQED}}^{(\text{int})} = \int_{\Omega} d^D x \sqrt{-g} \left(-i e g^{\alpha\beta} a_{\alpha} (\phi^* \partial_{\beta} \phi - \partial_{\beta} \phi^* \phi) + e^2 g^{\alpha\beta} a_{\alpha} a_{\beta} |\phi|^2 - \frac{g^4}{4} |\phi|^4 \right)$$

Scalar QED Interactions

Scalar-photon interactions:

$$S_{\text{sQED}}^{(\text{int})} = \int_{\Omega} d^D x \sqrt{-g} \left(-i e g^{\alpha\beta} a_{\alpha} (\phi^* \partial_{\beta} \phi - \partial_{\beta} \phi^* \phi) + e^2 g^{\alpha\beta} a_{\alpha} a_{\beta} |\phi|^2 - \frac{g^4}{4} |\phi|^4 \right)$$

Terms involved:

$$\mathcal{I}_{\{N\}}^{[\nu]} = \int_{-\infty}^{+\infty} du |\Delta u| u^{\nu} \prod_{i=1}^N \tilde{\Phi}_{\{k_{+(i)}, l_{(i)}, \vec{k}_{(i)}, r_{(i)}\}}(u)$$

$$\mathcal{J}_{\{N\}}^{[\nu]} = \int_{-\infty}^{+\infty} du |\Delta| |u|^{1+\nu} \prod_{i=1}^N \tilde{\Phi}_{\{k_{+(i)}, l_{(i)}, \vec{k}_{(i)}, r_{(i)}\}}(u)$$

Scalar QED Interactions

Scalar-photon interactions:

$$S_{\text{sQED}}^{(\text{int})} = \int_{\Omega} d^D x \sqrt{-g} \left(-i e g^{\alpha\beta} a_{\alpha} (\phi^* \partial_{\beta} \phi - \partial_{\beta} \phi^* \phi) + e^2 g^{\alpha\beta} a_{\alpha} a_{\beta} |\phi|^2 - \frac{g^4}{4} |\phi|^4 \right)$$

Terms involved:

$$\mathcal{I}_{\{N\}}^{[\nu]} = \int_{-\infty}^{+\infty} du |\Delta u| u^{\nu} \prod_{i=1}^N \tilde{\Phi}_{\{k_{+(i)}, l_{(i)}, \vec{k}_{(i)}, r_{(i)}\}}(u)$$

$$\mathcal{J}_{\{N\}}^{[\nu]} = \int_{-\infty}^{+\infty} du |\Delta| |u|^{1+\nu} \prod_{i=1}^N \tilde{\Phi}_{\{k_{+(i)}, l_{(i)}, \vec{k}_{(i)}, r_{(i)}\}}(u)$$

most terms **do not converge** due to **isolated zeros** ($l_{(*)} \equiv 0$) and cannot be recovered even with a **distributional interpretation** due to the term $\propto u^{-1}$ in the exponential

String and Field Theory

So far:

- field theory presents **divergences** (see sQED)
- obvious ways to regularise (Wilson lines, higher derivative couplings, etc.) **do not work**
- divergences are **not (only) gravitational**
- **vanishing volume** in phase space responsible for the divergence

String and Field Theory

So far:

- field theory presents **divergences** (see sQED)
- obvious ways to regularise (Wilson lines, higher derivative couplings, etc.) **do not work**
- divergences are **not (only) gravitational**
- **vanishing volume** in phase space responsible for the divergence

What about string theory?

String and Field Theory

So far:

- field theory presents **divergences** (see sQED)
- obvious ways to regularise (Wilson lines, higher derivative couplings, etc.) **do not work**
- divergences are **not (only) gravitational**
- **vanishing volume** in phase space responsible for the divergence

What about string theory?

Massive String States

$$V_M(x; k, S, \xi) =: \left(\frac{i}{\sqrt{2\alpha'}} \xi_\alpha \partial_x^2 X^\alpha(x, x) + \left(\frac{i}{\sqrt{2\alpha'}} \right)^2 S_{\alpha\beta} \partial_x X^\alpha(x, x) \partial_x X^\beta(x, x) \right) e^{ik \cdot X(x, x)}.$$

String and Field Theory

So far:

- field theory presents **divergences** (see sQED)
- obvious ways to regularise (Wilson lines, higher derivative couplings, etc.) **do not work**
- divergences are **not (only) gravitational**
- **vanishing volume** in phase space responsible for the divergence

What about string theory?

Massive String States

$$V_M(x; k, S, \xi) =: \left(\frac{i}{\sqrt{2\alpha'}} \xi_\alpha \partial_x^2 X^\alpha(x, x) + \left(\frac{i}{\sqrt{2\alpha'}} \right)^2 S_{\alpha\beta} \partial_x X^\alpha(x, x) \partial_x X^\beta(x, x) \right) e^{ik \cdot X(x, x)}.$$

*string theory cannot do **better than field theory** (EFT) if the latter **does not exist***

Resolution and Motivation

Introduce the generalised NBO:

$$\begin{cases} u &= x^- \\ z &= \frac{1}{2x^-} \left(\frac{x^2}{\Delta_2} + \frac{x^3}{\Delta_3} \right) \\ w &= \frac{1}{2x^-} \left(\frac{x^2}{\Delta_2} - \frac{x^3}{\Delta_3} \right) \\ v &= x^+ - \frac{1}{2x^-} \left((x^2)^2 + (x^3)^2 \right) \end{cases}$$

$$\Rightarrow \quad \kappa = -2\pi i (\Delta_2 J_{+2} + \Delta_3 J_{+3}) = 2\pi \partial_z$$

Resolution and Motivation

Introduce the **generalised NBO**:

$$\begin{cases} u &= x^- \\ z &= \frac{1}{2x^-} \left(\frac{x^2}{\Delta_2} + \frac{x^3}{\Delta_3} \right) \\ w &= \frac{1}{2x^-} \left(\frac{x^2}{\Delta_2} - \frac{x^3}{\Delta_3} \right) \\ v &= x^+ - \frac{1}{2x^-} \left((x^2)^2 + (x^3)^2 \right) \end{cases} \Rightarrow \kappa = -2\pi i (\Delta_2 J_{+2} + \Delta_3 J_{+3}) = 2\pi \partial_z$$

No isolated zeros $\Rightarrow$ distributional Interpretation

$$\tilde{\Phi}_{\{k_+, p, l, \vec{k}, r\}}(u) = \frac{1}{2\sqrt{(2\pi)^D |\Delta_2 \Delta_3 k_+|}} \frac{1}{|u|} e^{-i \left(\frac{1}{8k_+ u} \left[\frac{(l+p)^2}{\Delta_2^2} + \frac{(l-p)^2}{\Delta_3^2} \right] - \frac{\|\vec{k}\|^2 + r}{2k_+} u \right)}$$

On the Divergences and Their Nature

- divergences are present in sQED and **open string** sector
- singularities $\Rightarrow$ **massive states** are no longer spectators
- vanishing volume (**compact orbifold directions**) $\Rightarrow$ particles “cannot escape”
- **non compact** orbifold directions $\Rightarrow$ interpretation of **amplitudes as distributions**
- issue not restricted to NBO/GNBO but also BO, null brane, etc. (it is a **general issue** connected to the geometry of the underlying space)

On the Divergences and Their Nature

- divergences are present in sQED and **open string** sector
- singularities $\Rightarrow$ **massive states** are no longer spectators
- vanishing volume (**compact orbifold directions**) $\Rightarrow$ particles “cannot escape”
- **non compact** orbifold directions $\Rightarrow$ interpretation of **amplitudes as distributions**
- issue not restricted to NBO/GNBO but also BO, null brane, etc. (it is a **general issue** connected to the geometry of the underlying space)

*divergences are **hidden into EFT contact terms** and interactions with **string massive states**: gravity is not the only cause as the same problems are present also in gauge theories.*

[Arduino, RF, Pesando (2020)]

Contents

Conformal Symmetry and Geometry of the Worldsheet

Preliminary Concepts and Tools

D-branes Intersecting at Angles

Fermions and Point-like Defect CFT

Cosmological Backgrounds and Divergences

Orbifolds and Cosmological Toy Models

Null Boost Orbifold

Deep Learning the Geometry of String Theory

Machine Learning and Deep Learning

Machine Learning for String Theory

AI Implementations for Geometry and Strings

The Simplest Calabi–Yau

Focus on Calabi–Yau 3-folds:

$$h^{r,s} = \dim_{\mathbb{C}} H_{\bar{\partial}}^{r,s}(M, \mathbb{C}) \quad \Rightarrow \quad \begin{cases} h^{0,0} & = h^{3,0} = 1 \\ h^{r,0} & = 0 \quad \text{if } r \neq 3 \\ h^{r,s} & = h^{3-r,3-s} \\ h^{1,1}, h^{2,1} \in \mathbb{N} \end{cases}$$

The Simplest Calabi–Yau

Focus on Calabi–Yau 3-folds:

$$h^{r,s} = \dim_{\mathbb{C}} H_{\bar{\partial}}^{r,s}(M, \mathbb{C}) \quad \Rightarrow \quad \begin{cases} h^{0,0} & = h^{3,0} = 1 \\ h^{r,0} & = 0 \quad \text{if } r \neq 3 \\ h^{r,s} & = h^{3-r,3-s} \\ h^{1,1}, h^{2,1} & \in \mathbb{N} \end{cases}$$

Complete Intersection Calabi–Yau Manifolds

Intersection of hypersurfaces in

$$\mathcal{A} = \mathbb{P}^{n_1} \times \dots \times \mathbb{P}^{n_m}$$

where

$$\mathbb{P}^n: \quad \begin{cases} p_i(Z^0, \dots, Z^n) & = P_{i_1 \dots i_i} Z^{i_1} \dots Z^{i_i} = 0 \\ p_i(\lambda Z^0, \dots, \lambda Z^n) & = \lambda^i p_i(Z^0, \dots, Z^n) \end{cases}$$

[Green, Hübsch (1987); Hübsch (1992)]

Representation of the Output

CICY can be generalised to m projective spaces and k equations. The problem is thus mapped to:

$$\mathcal{R}: \quad \mathbb{N}^{m \times k} \quad \longrightarrow \quad \mathbb{N}$$

$$\left[\begin{array}{c|ccc} \mathbb{P}^{n_1} & a_1^1 & \dots & a_k^1 \\ \vdots & \vdots & \ddots & \vdots \\ \mathbb{P}^{n_m} & a_1^m & \dots & a_k^m \end{array} \right] \longrightarrow h^{1,1} \quad \text{or} \quad h^{2,1}$$

Representation of the Output

CICY can be generalised to m projective spaces and k equations. The problem is thus mapped to:

$$\mathcal{R}: \mathbb{N}^{m \times k} \longrightarrow \mathbb{N}$$

$$\left[\begin{array}{c|ccc} \mathbb{P}^{n_1} & a_1^1 & \dots & a_k^1 \\ \vdots & \vdots & \ddots & \vdots \\ \mathbb{P}^{n_m} & a_1^m & \dots & a_k^m \end{array} \right] \longrightarrow h^{1,1} \text{ or } h^{2,1}$$

Representation of the Output

CICY can be generalised to m projective spaces and k equations. The problem is thus mapped to:

$$\mathcal{R}: \mathbb{N}^{m \times k} \longrightarrow \mathbb{N}$$

$$\left[\begin{array}{c|ccc} \mathbb{P}^{n_1} & a_1^1 & \dots & a_k^1 \\ \vdots & \vdots & \ddots & \vdots \\ \mathbb{P}^{n_m} & a_1^m & \dots & a_k^m \end{array} \right] \longrightarrow h^{1,1} \text{ or } h^{2,1}$$

Machine Learning Approach

What is $\mathcal{R}$ in **machine learning** approach?

$$\mathcal{R}(M) \rightarrow \mathcal{R}_n(M; w) \rightarrow \hat{h}^{p,q} \quad \text{s.t.} \quad \exists n > M > 0 \quad | \quad \mathcal{L}_n(\hat{h}^{p,q}, h^{p,q}) < \epsilon \quad \forall \epsilon > 0$$

Machine Learning

- **optimisation problem** $\Rightarrow$ gradient descent (or similar)

Machine Learning



- **optimisation problem** $\Rightarrow$ gradient descent (or similar)
- use **various algorithms** and exploit **large datasets** (more training)

Machine Learning

- **optimisation problem** $\Rightarrow$ gradient descent (or similar)
- use **various algorithms** and exploit **large datasets** (more training)
- intersection of **computer science, mathematics and physics**

Machine Learning

- **optimisation problem** $\Rightarrow$ **gradient descent** (or similar)
- use **various algorithms** and exploit **large datasets** (more training)
- intersection of **computer science, mathematics and physics**
- provide in-depth **data analysis** of the datasets





[Green *et al.* (1987)]

Dataset

- 7890 CICY manifolds (full dataset)
- **dataset pruning**: no product spaces, no “very far” outliers (reduction of 0.49%)
- $h^{1,1} \in [1, 16]$ and $h^{2,1} \in [15, 86]$
- 80% training, 10% validation, 10% test
- choose **regression**, but evaluate using **accuracy** (round the result)

Dataset

- 7890 CICY manifolds (full dataset)
- **dataset pruning**: no product spaces, no “very far” outliers (reduction of 0.49%)
- $h^{1,1} \in [1, 16]$ and $h^{2,1} \in [15, 86]$
- 80% training, 10% validation, 10% test
- choose **regression**, but evaluate using **accuracy** (round the result)



Exploratory Data Analysis




exploratory data analysis → feature **selection** → Hodge numbers

[Ruehle (2020); Erbin, RF (2020)]


Exploratory Data Analysis

exploratory data analysis → feature **selection** → Hodge numbers

[Ruehle (2020); Erbin, RF (2020)]




[from scikit-learn.org]




Machine Learning

[from scikit-learn.org]



[from scikit-learn.org]



A Word on PCA




What is PCA for a $X \in \mathbb{R}^{n \times p}$?

- project data such that **variance is maximised**
- **eigenvectors** of XX^T or the **singular values** of X
- isolate **signal** from **background**
- ease the ML job of finding a better representation of the input

A Word on PCA


What is PCA for a $X \in \mathbb{R}^{n \times p}$?

- project data such that **variance is maximised**
- **eigenvectors** of XX^T or the **singular values** of X
- isolate **signal** from **background**
- ease the ML job of finding a better representation of the input




Machine Learning Results

Configuration Matrix Only




Machine Learning Results

Configuration Matrix Only



Best Training Set [Erbin, RF (2020)]



Artificial Intelligence and Neural Networks

- use **gradient descent** to optimise **weights**
- learn highly **non linear** representations of the input
- can be “large” to have enough parameters
- can be “deep” to learn **complicated functions**


Layers

fully connected: $\phi(a^{(i)} \{l\} \cdot W^{(l)} + b^{(l)} \mathbf{1})$

convolutional: $\phi(a^{(i)} \{l\} * W^{(l)} + b^{(l)} \mathbf{1})$

Non linearity ensured by:

$$\phi(z) = \text{ReLU}(z) = \max(0, z)$$




[rendition of the neural network in Bull et al. (2018)]

Convolutional Neural Networks

Why convolutional?

- retain **spacial awareness**
- smaller **no. of parameters**
($\approx 2 \times 10^5$ vs. $\approx 2 \times 10^6$)
- weights are **shared**
- CNNs isolate “**defining features**”
- find patterns as in **computer vision**



Convolutional Neural Networks


Why convolutional?

- retain **spacial awareness**
- smaller **no. of parameters**
($\approx 2 \times 10^5$ vs. $\approx 2 \times 10^6$)
- weights are **shared**
- CNNs isolate “**defining features**”
- find patterns as in **computer vision**

Convolutional Neural Networks

Why convolutional?

- retain **spacial awareness**
- smaller **no. of parameters**
($\approx 2 \times 10^5$ vs. $\approx 2 \times 10^6$)
- weights are **shared**
- CNNs isolate “**defining features**”
- find patterns as in **computer vision**



Inception Neural Networks


Recent development by deep learning research at Google led to:

- neural networks with better **generalisation properties**
- **smaller** networks (both parameters and depth)
- different **concurrent kernels**

Inception Neural Networks


Recent development by deep learning research at Google led to:

- neural networks with better **generalisation properties**
- **smaller** networks (both parameters and depth)
- different **concurrent kernels**






Deep Learning Results and Generalisation Properties

Best Training Set [Erbin, RF (2020)]



Deep Learning Results and Generalisation Properties

Best Training Set [Erbin, RF (2020)]



A Few Comments and Future Directions

Why deep learning in physics?

- o reliable **predictive method**

A Few Comments and Future Directions

Why deep learning in physics?

- reliable **predictive method** (provided good data analysis)
- reliable **source of inspiration**

A Few Comments and Future Directions

Why deep learning in physics?

- reliable **predictive method** (provided good data analysis)
- reliable **source of inspiration** (provided good data analysis)
- reliable **generalisation method**

A Few Comments and Future Directions

Why deep learning in physics?

- reliable **predictive method** (provided good data analysis)
- reliable **source of inspiration** (provided good data analysis)
- reliable **generalisation method** (provided good data analysis)
- **CNNs are powerful tools** (this is the *first time in physics!*)
- interdisciplinary approach = win-win situation!

A Few Comments and Future Directions

Why deep learning in physics?

- reliable **predictive method** (provided good data analysis)
- reliable **source of inspiration** (provided good data analysis)
- reliable **generalisation method** (provided good data analysis)
- **CNNs are powerful tools** (this is the *first time in physics!*)
- interdisciplinary approach = win-win situation!

What now?

- representation learning $\Rightarrow$ what is the best way to represent CICYs?
- study invariances $\Rightarrow$ invariances should not influence the result (graph representations?)
- higher dimensions $\Rightarrow$ what about CICY 4-folds?
- geometric deep learning $\Rightarrow$ explain the geometry of the “AI” behind deep learning!
- reinforcement learning $\Rightarrow$ give the rules, not the result!

The End?

- general framework for **D-branes at angles**
- alternative computations of **correlators of spin fields**
- strings and divergences in **time dependent orbifolds**
- string compactifications and **deep learning techniques**



THANK YOU